Michele Freppaz, Margherita Maggioni

**Università degli Studi di Torino
NatRisk - LNSA**

Snow cover duration/soil nutrient cycling

Snow cover duration/soil nutrient cycling

M. Freppaz et al. / Applied Soil Ecology 62 (2012) 131–141

Snow cover duration/soil nutrient cycling

Daily soil temperature amplitude

< 1-2 °C

Snow cover on the ground

Figure 7. A conceptual model of how snow cover controls subnival heterotrophic activity in alpine catchments at mid-latitudes. The duration of winter snow cover increases along the X axis and total over-winter heterotrophic activity and N leachate increase along the Y axes. The snow-pack controls on subnival microbial activity in turn exert control on the assimilation of inorganic N released from the snowpack and the export of NO_3^- to surface waters

Snow cover duration

211-240 giorni

2010	
Giorni	
91 - 120	241 - 270
121 - 150	271 - 300
151 - 180	301 - 330
181 - 210	331 - 360
211 - 240	Vannino
0 - 30	
31 - 60	
61 - 90

2010

Validazione con nivometri a terra

- 2000-2010
- 506 scene MODIS
- 9 stazioni della Rete ARPA Piemonte
- **Data di scomparsa della neve:** ad alta quota i nivometri sembrano sovrastimare, a bassa quota sottostimare

I nivometri e la fenologia delle specie erbacee

Plants as ecosystem engineers

Microtopografia:
i suoli a cuscinetti erbosi
(Earth Hummocks)

- Sollevamenti differenziali da gelo
- Cryoespulsione di clasti
- Accumulo di lettiera e materiale vegetale

Obiettivi

- Determinazione delle principali caratteristiche chimico-fisiche dei suoli a cuscinetti erbosi in corrispondenza di *hummocks* (“**up**”) e *interhummocks* (“**down**”)
- Valutazione delle caratteristiche della soluzione del suolo e andamento stagionale (primavera-estate-autunno)

Inquadramento geografico

- Quota: 2160 m.s.l.m.
- Comune di Torgnon, Loc. Tellinod
 - Esposizione Sud
- T media annua aria: 3.1 C°

Tesi Laurea Pintaldi, 2014 • Precipitazioni medie annue: 880 mm/anno

Inquadramento meteo-nivologico

- Altezza manto nevoso: 90-120 cm
- Durata media manto nevoso: 6 mesi
- Temperatura invernale suolo: $\sim 0\text{ C}^\circ$

Sito LTER di Tellinod

- Monitoraggio flussi CO₂
- Monitoraggio dinamiche stagionali e processi funzionali vegetazione
- Effetto dinamiche stagionali (durata innevamento e fusione) del manto nevoso

Inquadramento morfologico

Inquadramento vegetazionale

Zone UP:
Nardus Stricta

→
95-99%

Zone DOWN:
*Crocus vernus, Geum
montanum, Trifolium
alpinum, Arnica
montana, Polygonum
bistorta...*

→
45-75%

Campionamento del suolo

- Data campionamento: 8/11/2013
- Prelievo del *topsoil* (0-10 cm)
- 60 campioni (30 *up*-30 *down*)

Campionamento soluzione del suolo

- Lisimetri (prelievo 20 e 40 cm di profondità)
- Prelievi da giugno a dicembre 2013

N°PRELIEVO	DATA
1	06/06/2013
2	14/06/2013
3	26/06/2013
4	04/07/2013
5	24/07/2013
6	22/08/2013
7	08/11/2013
8	10/12/2013

Analisi di laboratorio

Materiali e
metodi

Tessitura

- Sabbiosa-franca (75% sabbia- 21% limo- 3.7 % argilla)
- Limo grossolano (0.05-0.02 mm)*

* Differenza significativa $p < 0.05$

pH, C_{org}

C/N

**Risultati
suolo**

Extr-DOC, DON, DIN

Risultati
suolo

Microtopografia

Pattern distribuzione manto nevoso

Data di fusione manto nevoso

Regime idrico e termico del suolo

Mineralizzazione SOM

**Risultati
suolo**

Frost as ecosystem engineer

Suoli con morfologia superficiale a *patterned ground* nel Parco Nazionale Gran Paradiso (Piata Lazin - Valle Soana)

D'Amico et al., submitted

- In ambienti caratterizzati da basse temperature, il *patterned ground* è una delle manifestazioni più spettacolari dell'azione del gelo.
- Il *patterned ground* è classificato sulla base di:
 - a) Forma geometrica (es. cerchi, poligoni, strisce)
 - b) Presenza o assenza di segregazione (sorted o unsorted)

Aconcagua, Argentina (32°45' S, 3428 m slm)

Alcuni esempi

Sorted
circles

Sorted
labyrinths

Sorted
stripes

Sorted
polygons

Kessler e Werner (2003)

Vorkuta (67°30' N; 64°10' E, 180 m slm), settembre 2004

PATTERN TYPE AND SLOPE ANGLE

▪ I processi responsabili dell'origine dei *sorted patterned ground* non sono ancora completamente noti.

Recentemente Kessler e Werner (2003) hanno sviluppato un modello, secondo il quale i cicli gelo/disgelo:

- 1) Inducono la separazione del materiale grossolano e la concentrazione del materiale fine in aree separate;
- 2) Favoriscono il trasporto del materiale grossolano lungo l'asse maggiore.

Area di Studio

- Valle Soana
- Parco Nazionale Gran Paradiso
- Piata Lazin

Area di Studio

- Plateau Sommitale (3028 m slm)
- Dislivello: 2000 m circa
- Gneiss occhiadino
- Temperatura media annua: -4°C
- Precipitazioni medie annue: 1600 mm

Area di Studio

D'Amico et al., submitted

Attività di campo ed analisi di laboratorio (1)

- Mappatura sorted circles
- Prelievo tospsoil in 2 sorted circles: pH, colore, Corg, Ntot, tessitura apparente, indice COLE

Attività di campo ed analisi di laboratorio (2)

- Misura temperatura suolo: ottobre 2007-ottobre 2008 (data loggers UTL-1)
- Stazione nivometeorologica Lago Valsoera (SMI)

Mappatura sorted circles

- Numero sorted circles/ha:
233
- Diametro sorted circles: 0.5-
5m
- Dimensione clasti sui bordi:
5-25 cm
- H bordi: 6-12 cm

Piata Lazin (3085 m slm)

BTS: -3.9 °C

Probabilità presenza di permafrost: ALTA

Col d'Olen 2901 m slm

BTS: $-0.5\text{ }^{\circ}\text{C}$

Probabilità presenza di permafrost: BASSA

Ghiaia grossolana (20-75 mm), di forma angolare

Sostanza organica: $0.19 - 0.58 \text{ g}100\text{g}^{-1}$

▪ La presenza di vegetazione solo sui bordi esterni evidenzia come il fenomeno non sia più così espresso.

▪ In accordo con numerosi studi la presenza di specie vegetali indica condizioni relativamente stabili.

- La presenza di sorted circles più piccoli (5-10 cm) all'interno delle figure di maggiori dimensioni evidenzia però che i processi sono almeno in parte attivi.

- **Nel sito oggetto di studio la presenza del permafrost è probabile.**
- **L'organizzazione spaziale del materiale fine e del materiale grossolano determina una discontinuità nella distribuzione delle caratteristiche dei suoli e delle piante.**
- **Il processo di segregazione del materiale sembra ancora parzialmente attivo, con la formazione di forme “miniaturizzate” nelle porzioni centrali dei cerchi più grandi, che invece sembrano parzialmente stabilizzati.**

Snow gliding and loading under two different forest stands: a case study in the north-western Italian Alps

2 STUDY PLOTS

Larix decidua

Picea Abies

The climate of the area is continental humid subarctic, with a mean annual air temperature equal to + 4 °C and a mean annual precipitation equal to 718 mm. The mean annual snow depth is 175 cm (historical dataset from 1913 to 2001, Mercalli et al. 2003).

Two adjacent plots (40x40 m), located at the same elevation and aspect, were chosen (Fig. 1): a) one within a larch stand (Larch plot) and b) one within a spruce stand (Spruce plot). By counting all trees in each plot (with stem diameter > 5 cm) the average stem density was determined equal to 300 stems/ha for both plots. The crown canopy, measured as crown projection, amounted to 60% for the Larch plot and 65% for the Spruce plot. The average stem diameter was 40 cm and most of the trees were included in the range 30–40 cm (10 cm interval). The natural regeneration was generally scarce.

Physical snow properties

Table 1. Physical snow properties - according to Fierz et al. (2009) - measured in the field during winter 2009-2010 (8 snow profiles) and 2010–2011 (3 snow profiles), under Larch and Spruce plots.

Parameter	2009–2010		2010–2011	
	Larch	Spruce	Larch	Spruce
Maximum value of snow depth (cm)	60	66	28	22
Dominant grain type	FCso, MFpc	FCso, DHcp	RGsr	RGsr
Maximum grain size (mm)	2.5	4	1.5	1.5
Mean hand hardness Index	3	4	3	2
Mean snow density (kg·m ⁻³)	277	229	220	270
Dominant grain type in the basal layer	FCso, DHcp	DHcp	FCso	FCso
Mean snow/soil interface temperature (°C)	-3.8	-2.1	-1.1	-1.8
Date of snowpack isothermal condition	March 11 rd	March 19 rd	March 14 rd	March 20 rd

Note: FCso means solid faceted particles; MFpc means rounded polycrystals; DHcp means hollow cups; RGsr means small rounded particles.

Snow pressure on trees

Swiss guidelines

Margreth et al. (2007) applied the Swiss Guidelines with some improvements in order to calculate snow pressure on cableway masts. The snow pressure was calculated with the following equation (Eq.1):

$$S'_N = \rho g (H_S^2 / 2) KN \quad (1)$$

where: S'_N is the snow pressure component in the line of slope per meter run of the supporting surface along the contour line [$kN \cdot m^{-1}$], ρ is the density of the snow cover [$t \cdot m^{-3}$], g is the gravitational acceleration [ms^{-2}], H_S is the vertical snow depth [m]; K is the creep factor dependent on the slope inclination ψ and the density ρ , and N is the glide factor.

Mc Clung and Larsen (1989)

$$\sigma = \{ [2 / (1 - \nu)] [(d' / D) + (L / H_S)] \} 0.5 \rho g D \sin \psi + 0.5 [\nu / (1 - \nu)] \rho g D \cos \psi$$

where: σ is the maximal principal stress of the snow on a structure [Nm^{-2}], ν is the Poisson coefficient deduced from Salm et al. (1971). We used $\nu = 0.1$ that corresponds to a mean snow density of $250 \text{ kg} \cdot \text{m}^{-3}$, equal to the density we measured in our plots; L/H_S is a dimensionless creep parameter calculated by McClung and Larsen (1989) equal to $0.27 + (\nu/12)$, D is the snow thickness, ρ is the snow density measured in the field [$kg \cdot m^{-3}$], g is the gravitational acceleration [$m \cdot s^{-2}$], ψ is the slope angle, and d' is the stagnation depth.

The stagnation depth is a fundamental parameter to formulate the boundary condition at the snow/soil interface. We calculated the stagnation depth d' from the measured snow gliding rates using Equation 5 (McClung 1974):

$$d' = [\mu V_g] / [2(1 - \nu) \rho g D \sin \psi]$$

where: μ is the viscosity, equal to 10^{11} Pa for snow, V_g is the snow gliding velocity [mm/day]. According to Hoeller (2009) we used the maximum daily snow gliding rate recorded during winters 2009–2010 and 2010–2011; ν is the Poisson coefficient.

Fig. 3 Snow depth and air temperature measured by the automatic snow and weather station (a); T_i (b) M_i (c) and snow gliding (d) in each plot recorded in both winters (2009-2010 and 2010-2011) when the study site was snow-covered.

From our results, the snow/soil interface moisture significantly influenced the snow gliding processes. In particular, soil moisture at the snow/soil interface was related to the snow/soil interface temperature and both parameters (M_i and T_i) were strongly regulated by air temperature.

The volumetric moisture increase of a relatively dry snow/soil interface (10%–20%), seemed to be the most important factor influencing the snow gliding rates: in particular snow gliding was observed with wetting rates higher than 1.2%/day.

Modelling soil removal from snow avalanches: A case study in the North-Western Italian Alps

Confortola et al., 2012

Load: 25 t/ha

Superficial deposit: 1.8 mm

Table 1
Main features of the avalanche events.

Variable	Legend	Event		
		1	2	3
V_R [m ³]	Volume at deposition	$7 \cdot 10^4$	$6.3 \cdot 10^4$	$2.4 \cdot 10^3$
ρ_{in} [kgm ⁻³]	Snow density at first survey	580	515	670
ρ_{fin} [kgm ⁻³]	Snow density at last survey	630	480	-
S_{in} [m ²]	Avalanche deposit surface at first survey	$6.1 \cdot 10^4$	$2.9 \cdot 10^4$	$2.4 \cdot 10^4$
S_{fin} [m ²]	Avalanche deposit surface at last survey	315	$7.3 \cdot 10^3$	$2.4 \cdot 10^4$
C_s [kgm ⁻³]	Sediment concentration last survey	241	37	10
M_s [kg]	Mass of sediments	$7.6 \cdot 10^3$	$2.7 \cdot 10^4$	$2.3 \cdot 10^3$
V_0 [m ³]	Estimated avalanche volume at release	$1.6 \cdot 10^5$	$1.4 \cdot 10^5$	$2.4 \cdot 10^4$
H_0 [m]	Estimated avalanche depth at release	1.54	1.40	1.04
S_0 [m ²]	Estimated avalanche surface at release	$1 \cdot 10^5$	$1 \cdot 10^5$	$2.3 \cdot 10^4$
A_0 [m asl]	Average altitude at release	2652	2652	2622
A_R [m asl]	Altitude end mark	1058	1058	1098
L_R [m]	Run out distance	3532	3532	3038
A_S [m asl]	Snow line altitude	1980	1900	2000

AVALANCHE MODEL

$$MgA = MgRL, \quad (1)$$

with M [kg] avalanche body mass, A [m] vertical jump, L [m] runout length, g gravity [ms^{-2}], and R [.] flow resistance factor, *i.e.* energy dissipation.

The conservation equation of total energy E_{tot} [Joule] between two sections, namely Section 1 and Section 2 along the avalanche track is cast as

$$E_{tot}(1) = E_{tot}(2) + E_{dis}(1, 2), \quad (2)$$

with suffix *dis* indicating along track energy dissipation. The total energy can be split into potential and kinetic energy as

$$E_{tot} = E_{pot} + E_{kin} = MgA + \frac{MU^2}{2} \quad (3)$$

with U (vertically averaged) flow velocity. Merging of Eqs. (2) and (3) yields

$$MgA_1 + \frac{MU_1^2}{2} = MgA_2 + \frac{MU_2^2}{2} + E_{dis}. \quad (4)$$

The energy dissipation may be expressed using a Voellmy like approach (Voellmy, 1955) as

$$E_{dis} = MgL \cos \varphi \mu + \frac{MgLU^2}{\xi H} + MgL \cos \varphi \lambda \frac{\partial H}{\partial x}, \quad (5)$$

SOIL EROSION MODEL

Here we model ASR according to two different approaches, both based upon a hydraulic analogy, *i.e.* using classical approaches to soil removal and solid load in rivers (e.g. Graf, 1971; Yalin, 1982), namely i) excess of shear stress, ES, and ii) critical velocity, CV.

Confortola et al., 2012

SNOW GLIDING PARAMETERS:

snow depth, density, gliding factor (vegetation, topography, etc.,)

AVALANCHE PARAMETERS:

fracture depth (DH3gg), friction parameters, erodible snow cover,

→ **MODEL INPUTS**

CLIMATE CHANGE

???